

CONTINUOUS IMAGES OF CANTOR'S TERNARY SET

F. DREHER* AND T. SAMUEL†

ABSTRACT. Cantor's ternary set was initially discovered in 1874 by Henry Smith and has a number of remarkable and deep properties. Through its discovery, Georg Cantor and others helped lay the foundations of modern point-set topology. Among the many properties of Cantor's ternary set is the property that any compact metric space is the continuous image of Cantor's ternary set. This was first proved in 1927, with a proof occurring in the second addition of Felix Hausdorff's *Mengenlehre* and by Pavel Sergeyevich Alexandroff's article titled *Über stetige Abbildungen kompakter Räume*. Thus, a natural question to ask is, is it true that any compact topological space is the continuous image of the Cantor set? The answer to this question is no, since there exists a plethora of compact topological spaces whose cardinality exceeds that of the Cantor set. With this in mind, we are immediately led to the question, is it true that any compact topological space with the same cardinality as the continuum (i.e. that of Cantor's ternary set) is the continuous image of Cantor's ternary set? This question is precisely what we address in this note.

1. INTRODUCTION.

Through the efforts of Benoit B. Mandelbrot, his colleagues, co-authors and friends, fractals have been brought to the attention of a wider public. The simplest and probably the most important such set is Cantor's ternary set (also referred to as the middle third Cantor set), first introduced by Henry Smith [9] and subsequently and independently by Georg Cantor [2]. Cantor's ternary set is generated by the simple recipe of dividing the unit interval $[0, 1]$ into three parts, removing the open middle interval, and then continuing the process so that at each stage, each remaining subinterval is similarly subdivided into three and the middle open interval removed. Continuing this process *ad infinitum* one obtains a non-empty set consisting of an infinite number of points (see Figure 1).

Before formally introducing Cantor's ternary set, let us take a little time to discuss its history; our main sources for this are [4, 5]. The initial discovery of Cantor's ternary set was motivated by the theory of integration. For a while it was thought that a function on the real line would be Riemann-integrable if, and only if, it were point-wise discontinuous, with discontinuities occurring on a nowhere dense set. (A closed set is said to be nowhere dense if the largest open subset is the empty-set.) Moreover, it was believed that sets of the form $\{1/2^n : n \in \mathbb{N}\}$ were a prototype for all nowhere dense sets of the real line; specifically it was thought that all nowhere dense sets have zero outer Jordan content. Roughly speaking, this means that the set can be covered by finitely many intervals of arbitrarily small total length. However, in 1875 Henry Smith [9] proved the existence of nowhere dense sets with positive outer Jordan content. In fact he gave a specific recipe for constructing such sets, which nowadays are called *Smith-Cantor sets* or *fat-Cantor sets*. Similar to the construction of Cantor's ternary set, given in our opening paragraph, a Smith-Cantor set is constructed by removing certain open intervals from the unit interval. A particular example of such a set is built via the following procedure; one begins by removing the middle open interval centered about the point $1/2$ of length $1/2^2$ from the closed unit interval, so that the remaining set is the closed set $[0, 3/8] \cup [5/8, 1]$. The following steps consist of removing subintervals

Date: March 18, 2013.

* fdreher@uni-bremen.de - Fachbereich Mathematik, Universität Bremen, 28359 Bremen, Germany.

† tony@uni-bremen.de - Fachbereich Mathematik, Universität Bremen, 28359 Bremen, Germany.

of width $1/2^{2n}$ from the middle of each of the 2^{n-1} remaining intervals. Thus, for the second step, the open intervals $(5/32, 7/32)$ and $(25/32, 27/32)$ are removed from the closed set $[0, 3/8] \cup [5/8, 1]$. Continuing this process *ad infinitum*, the remaining set is called a Smith-Cantor set. This particular set is nowhere dense in the closed unit interval, but has outer Jordan content $1/2$.




FIGURE 1. The first six approximations to Cantor's ternary set.

A decade later, Georg Cantor, through his work on the point-wise convergence of trigonometric series rediscovered the set that is now referred to as Cantor's ternary set; specifically Georg Cantor was working on the question, if for all real numbers x , except those in some set P , we have that

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx)) = 0,$$

must all the coefficients a_n and b_n be zero? This led him to study the point-set topology of the real line and in a footnote in the fifth of a series of papers titled *Über unendliche, lineare Punktmannigfaltigkeiten*, Georg Cantor presents the set which we now refer to as Cantor's ternary set. Moreover, interestingly, it is believed that these articles were a precursor to Cantor's theory of transfinite numbers.

We have given the standard geometric definition of Cantor's ternary set in our opening paragraph, let us now formally define Cantor's ternary set in arithmetic terms.

Definition 1.1. The set

$$C := \left\{ \sum_{n \in \mathbb{N}} \frac{2 \cdot \omega_n}{3^n} : \omega_n \in \{0, 1\} \text{ for all } n \in \mathbb{N} \right\}$$

is called *Cantor's ternary set*.

Here and in the sequel we always let C denote Cantor's ternary set equipped with the subspace topology (Definition 2.4) induced by the Euclidean norm of $\mathbb{R}$. Below we list some of the remarkable and deep properties of Cantor's ternary set adhered to in the abstract; for the reader who is not familiar with terminology from point-set topology, we include, in Section 2, basic definitions from point-set topology. For a proof of Properties (2) to (9) we refer the reader to [3] or [8, Counterexample 29]; Property (1) is discussed in Section 3; for a proof of Properties (10), (11) and (12), the definition of the Lebesgue measure, Hausdorff dimension and that of a self-similar set, we refer the reader to [3].

- (1) Any compact metric space (Definition 2.1 and 2.9) is the continuous image of C .
- (2) The set C is the infinite product of locally connected spaces (Definition 2.6), but is not locally connected itself.
- (3) The set C is totally disconnected (Definition 2.7).

- (4) The set C is perfect (Definition 2.8).
- (5) The set C is compact (Definition 2.9).
- (6) The set C is nowhere dense in the closed unit interval $[0, 1]$.
- (7) The set C is Hausdorff (Definition 2.10).
- (8) The set C is normal (Definition 2.11).
- (9) The cardinality of C is equal to that of the continuum.
- (10) The one dimensional Lebesgue measure and outer Jordan content of C are both equal to zero.
- (11) The Hausdorff dimension of C is equal to $\log(2)/\log(3)$.
- (12) The set C is a self-similar set.

In this note we are interested in whether Property (1), more commonly known as the Hausdorff–Alexandroff Theorem, can be strengthened. What is not remarked, in the current literature, as far as we are aware, is the importance of the space being a metric space. Clearly, since there exists a plethora of compact topological spaces whose cardinality exceeds that of the Cantor set, the question we endeavour to answer is:

Does there exist a compact topological space with cardinality not exceeding that of the continuum (i.e. that of C) that is not the continuous image of C ?

The problem in proving or dis-proving this result is that one needs to consider spaces which are quite far away from being Hausdorff, since any compact, Hausdorff space which is second countable (that is, a space whose topology can be generated, see Definition 2.3, by a countable collection of sets) is metrisable. Moreover, out of those spaces which are non-Hausdorff, we need to exclude those which are the quotient or factor of a compact metric space. It turns out that the answer to this question is in the affirmative.

Theorem 1.1. *There exists a compact topological space (T, τ) with cardinality not exceeding that of the continuum which is not the continuous image of C when equipped with the subspace topology induced by the Euclidean norm of the real line $\mathbb{R}$.*

We will prove this result in Section 4. In doing so we will need an auxiliary result, Lemma 4.1, in the proof of this result we also rectify an error in [8, Counterexample 99]. However, before doing so, we first, in Section 2, present basic definitions from point set topology. Then, for completeness, in Section 3 we present an outline of the proof of the Hausdorff–Alexandroff Theorem (Property (1)).

2. DEFINITIONS AND EARLY HISTORY OF POINT-SET TOPOLOGY.

The definition of a topological space, that is now standard, was a long time in the making and generalises the notion of a metric space. Georg Cantor, Camille Jordan and Giuseppe Peano shaped the beginning of point-set topology in the late nineteenth century, which then developed further during the first decades of the twentieth century through the efforts of Adolf Hurwitz, Arthur Schoenflies, Henri Poincaré, Hans Hahn and David Hilbert, to name but a few.

2.1. The early history. Point-set topology is the study of the intrinsic properties of surfaces that are independent of distance. The classic example of this is that, from a topological point of view, the surface of a doughnut (or torus) and that of a coffee cup are the ‘same’, that is, one such object can be continuously deformed, without ripping, glueing or punching a hole, into the other. But why is point-set topology an important topic to understand? We believe that a brief historical recap of the beginnings of point-set topology should help a beginner grasp and become interested in this area, which is often remarked as being inaccessible. Our main source is [7].

Two very important results in analysis cannot, in modern mathematical terms, be discussed without the notion of a limit point. These two results are:

- (a) the Bolzano–Weierstraß Theorem, which states that every infinite set of points in a bounded region of $\mathbb{R}^n$ possesses at least one limit point, and
- (b) the existence of a function which is both nowhere differentiable and everywhere continuous, such a function is referred to as a *Weierstraß function*.

Thus, in the work of Bernardus Bolzano and Karl Weierstraß (early to mid nineteenth century) there are places where one can arguably see the beginnings of point-set topology. Georg Cantor’s series of papers titled *Über unendliche, lineare Punktmannigfaltigkeiten* addresses precisely this point and much more. In particular, how can one define a limit point. This problem is solved by introducing the derived set, a purely point-set notion. Moreover, in these articles Georg Cantor gives a detailed and meticulous construction of the real numbers and in doing so made it possible to build a rigorous foundation of the notion ‘nearness without distance’. These initial ideas place point-set topology within the larger mathematical world.

Camille Jordan and Giuseppe Peano were among the first to take Georg Cantor’s ideas further, and although they worked separately on problems of measure theory and integration, they saw the importance of Georg Cantor’s work and the consequences it had on the topic of measure theory and integration. Moreover, both Camille Jordan and Giuseppe Peano developed Georg Cantor’s work further. Within the topic of topology Camille Jordan is probably most famous for his celebrated theorem on closed curves in the plane and Giuseppe Peano for his astounding discovery/construction of a space filling curve, a curve which covers all points of a square (see [10] for an introduction to the topic of space filling curves).

2.2. Basic definitions. Let us now begin by defining various basic concepts, which we will use and have already mentioned above. For a good introduction to point-set topology and proofs of results mentioned but not proven, we refer the reader to [11, 12]. A metric space is a set of points whose only structure is a notion of distance.

Definition 2.1 (Metric space). A *metric space* is a set X of points together with a distance function $d : X \times X \rightarrow \mathbb{R}$ such that, for all $x, y, z \in X$:

- (Positive): $d(x, y) \geq 0$ and $d(x, y) = 0$ if, and only if, $x = y$,
- (Symmetric): $d(x, y) = d(y, x)$, and
- (Triangle inequality): $d(x, z) \leq d(x, y) + d(y, z)$.

Closely related to metric spaces are topological spaces, which are equipped with a notion of closeness without the need for a distance function. In the topological approach, the notion of an open set is used to ‘measure’ closeness. In particular two points are said to be ‘close’ if they share in common ‘most’ of their open neighbourhoods.

Definition 2.2 (Topological space). A *topological space* is a set of points X together with a collection of subsets τ of X , which are by definition the *open* subsets of X , such that:

- (a) any arbitrary union of open sets is open,
- (b) any finite intersection of open sets is open, and
- (c) the set X and the empty set $\emptyset$ are both open.

The collection of open sets τ is often referred to as the *topology*. An open set U containing a point $x \in X$ is called an *open neighbourhood* of x . A set $A \subseteq X$ whose complement A^c is open is called a *closed* set.

Definition 2.3 (Generating set). A *generating set* for a topology τ is a collection of sets $\gamma \subseteq \tau$ such that any set belonging to τ can be written as the finite intersection of arbitrary unions of sets from γ .

Definition 2.4 (Subspace topology). All subsets $A \subset X$ inherit a topology from (X, τ) , defined by intersecting each element of τ with A .

Topologies may be compared to each other. If all open sets in a topology τ_1 are also open in τ_2 , then τ_2 is said to be *finer* than τ_1 , while τ_1 is said to be *coarser* than τ_2 . The coarsest topology is the *trivial topology* given by the collection $\{\emptyset, X\}$, while the finest topology is the *discrete topology*, in which every subset of X is defined to be open. Two topologies are said to be *equivalent* if τ_1 is finer than τ_2 and vice versa.

In a metric space or normed space, the generating set given by the set of open balls forms the topology called the *metric topology* sometimes referred to as the topology induced by the metric, or respectively, the norm. Note that the set of open balls itself is not a topology, since unions of open balls are open but may not themselves be open balls. Further, A topological space is said to be *metrisable* if there exists a metric on the space that induces an equivalent topology.

2.2.1. Functions between topological spaces. In topology, continuity is expressed in terms of open sets. A *continuous function* $f : X \rightarrow Y$ between topological spaces (X, τ_1) and (Y, τ_2) is a function such that $f^{-1}(U)$ is open whenever $U \subseteq Y$ is open. A function is continuous at a point $x \in X$ if it is continuous on an open neighbourhood of x . This definition coincides with the ϵ - δ definition of a continuous function between metric spaces. Note the directionality of this definition, in particular one requires the inverse image of open sets to be open, rather than the forward image. There is another name for when the forward image of an open set is open, namely, a function $f : X \rightarrow Y$ is said to be *open (closed)* if $f(A)$ is open (closed) whenever A is open (closed).

2.2.2. Properties of topological spaces. The notion of open and closed sets also permits a description of when a set consists of a 'single piece', instead of several smaller pieces.

Definition 2.5 (Connected). A space X which is not the disjoint union of two non-empty open subsets is called *connected*.

Definition 2.6 (Locally connected). Let (X, τ) be a topological space, and let x be a point of X . We call X *locally connected at x* if for every open set V containing x there exists a connected, open set U with $x \in U \subseteq V$. The space X is said to be *locally connected* if it is locally connected at all $x \in X$.

Definition 2.7 (Totally disconnected). Let (X, τ) be a topological space. The maximal connected subset of X containing a point x is called the *connected component* of x . The topological space (X, τ) is called *totally disconnected* if the only connected components of X are single points.

Definition 2.8 (Perfect). Let $A \subseteq X$ be a subset of a topological space (X, τ) . A point x in X is a *limit point* of A if every open neighbourhood of x contains at least one point of A different from x itself. The set A is called *perfect* if every point $x \in A$ is a limit point of A .

Since topological spaces have no notion of distance, they do not really have a notion of size. In some sense, however, the notion of compactness in essence encapsulates this concept. In particular, compact spaces allow a transformation from the infinite to the finite.

Definition 2.9 (Compact). Given a subset $A \subseteq X$ of a topological space (X, τ) , an *open cover* of A is a collection of open sets whose union contains A . A *subcover* is a sub-collection of an open cover whose union still contains A . We call a subset A of X *compact* if every open cover has a finite subcover.

It is not difficult to show that if $f : X \rightarrow Y$ is a continuous function between two topological spaces (X, τ_1) and (Y, τ_2) , and if A is a compact subset of X , then $f(A)$ is also compact. Moreover, one can show that a subset A of n -dimensional Euclidean space equipped with the topology induced by the Euclidean norm, is compact if, and only if, it is closed and bounded, that is, there exists a constant $\kappa > 0$ such that for all $a \in A$, the norm of a is less than κ . This is known as the Heine–Borel Theorem.

Another result which will be important in the sequel is that any closed subset of a compact topological space is compact.

Before ending this section, we state a key result in topology, which we adhered to in the introduction, that describes when a topological space is metrisable.

Theorem 2.1. *A compact topological space is second countable if and only if it is metrisable.*

2.2.3. Separation properties of topological spaces. In topology, separation indicates the ability of open sets to distinguish points. The separation axioms classify a topological space X based on how easy it is to separate points into different subsets of X . In this article we will be mainly concerned with the following two separation axioms.

Definition 2.10 (Hausdorff). A topological space (X, τ) is called *Hausdorff* if given distinct points $x, y \in X$, there are disjoint open sets $U \ni x$ and $V \ni y$.

Definition 2.11 (Normal). A topological space (X, τ) is called *normal* if given disjoint closed sets $F, G \subset X$, there are disjoint open sets $U \supseteq F$ and $V \supseteq G$.

Remark. Any metrisable topological space is Hausdorff and normal. In particular, Cantor's ternary set is Hausdorff and normal. (Recall that we assume throughout this article that Cantor's ternary set is equipped with the subspace topology induced by the Euclidean norm.)

3. THE HAUSDORFF-ALEXANDROFF THEOREM.

In this section we present a sketch of the proof of the Hausdorff–Alexandroff Theorem which can be found in most modern books on point set topology, for instance [12, Theorem 30.7] or [10, Theorem 6.6]. The theorem was first proved in 1927, in the second addition of Felix Hausdorff's *Mengenlehre* and independently by Pavel Sergeyevich Alexandroff [1]. This result is not only interesting in its own right, but allowed Hans Hahn to provide a simpler proof of his 1914 theorem [6] which was also independently proven by Stefan Mazurkiewicz. The statement of this result, which is now known as the Hahn–Mazurkiewicz Theorem, is that a Hausdorff topological space is the continuous image of the closed unit interval if, and only if, it is a compact, connected, locally connected metric space. We refer the reader to [12, Theorem 31.5] for a complete proof of the statement, with the remark that the difficulty lies in showing the backward implication.

Theorem 3.1 (Hausdorff–Alexandroff Theorem). *Any compact metric space K is the continuous image of Cantor's ternary set C .*

In order to outline a proof of this result, we require one further definition, and that is the closure of a subset of a topological space. An important property of the closure of a set, which will be used, is that it is closed with respect to the topology it is taken with.

Definition 3.1 (Closure of a set.). Given a topological space (T, τ) and a set $U \subset T$, we define the closure of the set U with respect to τ to be the set U together with all of its limit points under the given topology τ . We denote this latter set by $\overline{U}$.

Sketch of Proof. The following is based on [10] and follows the same basic steps as Felix Hausdorff's original proof. Pavel Sergeyevich Alexandroff's proof, on the other hand, differs significantly and, interestingly, uses continued fractions.

From the arithmetic description of Cantor's ternary set C , it is easily verified that each point in C corresponds to a 0-1 sequence. If we can devise a 0-1 coding for the points in the compact metric space K , we can easily define a mapping $f : C \rightarrow K$ which then only needs to be checked for continuity.

We cover $K_0 := K$ by open balls of radius $1/2^0 = 1$. Since K is compact we can choose a finite subcover. If necessary we can include more sets than required or even include some sets multiple times to obtain a finite cover (U_{i_1}) consisting of 2^{n_1} sets, for some natural

number n_1 . For each set U_{i_1} of this subcover we define a new compact set $K_{i_1} := \overline{U_{i_1}} \cap K_0$ and repeat the procedure for each of the K_{i_1} with open balls of radius $1/2^1$. Since these are only finitely many sets we can cover each K_{i_1} by 2^{n_2} sets, for some natural number n_2 independent of i_1 . Continuing this process and halving the radius of the open balls in each step we arrive at nested sequences $K = K_0 \supset K_{i_1} \supset K_{i_1, i_2} \supset \dots$ with $i_k \in \{0, 1, \dots, 2^{n_k-1}\}$ for each natural number k . Since K is compact and the diameter of the sets decreases with each step by a factor of $1/2$, the intersection of each such nested sequence is non-empty and consists of exactly one point (this is known as Cantor's Intersection Theorem). Also, each point of K is thusly represented by at least one of these sequences because at each finite level the sets form a cover of K .

We have chosen 2^{n_k} sets at each step and therefore we can transform the (i_k) -coding into a full binary coding. Now we can define a mapping $f : C \rightarrow K$ that maps the binary-coded points of the Cantor set surjectively (but usually not injectively) onto the now also binary-coded points of K . This mapping is continuous because the binary codings of the entries of a converging sequence $x_k \rightarrow x$ in C coincide in an increasing number of places with the coding of x and thus their images $f(x_k)$ are contained in a nested sequence of compact sets with $f(x)$ as the one point in the intersection. $\square$

Remark. The continuity argument relies on the fact that convergence in C with regard to the metric directly corresponds to convergence in the binary expansion. This only works because of the unique binary coding in a ternary environment that Cantor's ternary set exhibits; one cannot use the same argument with the usual non-unique binary coding of the elements in the interval $[0, 1]$.

4. PROOF OF THEOREM 1.1.

The main idea is to prove Theorem 1.1 by contradiction. In particular to choose a specific non-Hausdorff space and show that if there exists a continuous map from the Cantor set into this space, then the continuous map must push-forward the Hausdorff property of Cantor's ternary set, which will be a contradiction to how the target space was originally chosen. Let us first describe the non-Hausdorff space which we will take as our target space, after which we will present the proof of Theorem 1.1.

Lemma 4.1. *There exists a countable space T with a topology τ which has the following properties:*

- (a) (T, τ) is compact,
- (b) (T, τ) is non-Hausdorff, and
- (c) every compact subset of T is closed with respect to τ .

Proof. This proof is inspired by [8, Counterexample 99]. We define

$$T := (\mathbb{N} \times \mathbb{N}) \cup \{x, y\},$$

namely the Cartesian product of the set of natural numbers $\mathbb{N}$ with itself unioned with two distinct arbitrary points x and y . We equip the set T with the topology τ generated by the sets described in (i), (ii) and (iii) given directly below.

- (i) The sets of the form $\{(m, n)\}$, for each $(m, n) \in \mathbb{N} \times \mathbb{N}$.
- (ii) The sets $T \setminus A$, where $A \subset (\mathbb{N} \times \mathbb{N}) \cup \{y\}$ contains y and is such that the cardinality of the set $A \cap \{(m, n) : n \in \mathbb{N}\}$ is finite for all $m \in \mathbb{N}$, that is, the set A contains at most finitely many points on each row. These sets are the open neighbourhoods of x .
- (iii) The sets $T \setminus B$, where $B \subset (\mathbb{N} \times \mathbb{N}) \cup \{x\}$ contains x and is such that there exists an $M \in \mathbb{N}$, so that if $(m, n) \in A \cap (\mathbb{N} \times \mathbb{N})$, then $m \leq M$, that is B contains only points from at most finitely many rows. These sets are the open neighbourhoods of y .

Property (a) follows from the observation that any open cover of T contains at least one open neighbourhood $T \setminus A$ of x and one open neighbourhood $T \setminus B$ of y with A and

B as given above. The points not already contained in these two open sets are contained in $T \setminus ((T \setminus A) \cup (T \setminus B)) = A \cap B$ which, by construction, is a finite set. In this way a finite subcover can be chosen and hence the topological space (T, τ) is compact.

To see why T has Property (b), consider open neighbourhoods of x and y . An open neighbourhood of x contains countably infinitely many points on each row of the lattice $\mathbb{N} \times \mathbb{N}$; an open neighbourhood of y contains countably infinitely many full rows. It follows that there are no disjoint open neighbourhoods $U \ni x$ and $V \ni y$ and thus T is non-Hausdorff.

We use contraposition to prove Property (c). Suppose that $E \subset T$ is not closed. Note that we may assume that E is a strict subset of T since T itself is closed by the fact that $\emptyset \in \tau$. By construction of the topology, a set that is not closed cannot contain both x and y . Also, there needs to be at least one point in the closure $\overline{E}$ of E but not already in E ; this has to be one of the points x or y , because singletons $\{(m, n)\}$ which are subsets of the lattice $\mathbb{N} \times \mathbb{N}$ are open and thus the point (m, n) cannot be a limit point of E . We shall now check both cases, that is, (i) if $x \in \overline{E} \setminus E$ and (ii) if $y \in \overline{E} \setminus E$.

- (i) If $x \in \overline{E} \setminus E$, then every open neighbourhood of x has a non-empty intersection with E . It follows that there is at least one row in $\mathbb{N} \times \mathbb{N}$ that shares infinitely many points with E . Denote this row by B . Then the open cover $\{T \setminus B\} \cup \{b\} : b \in B \cap E\}$ of E cannot be reduced to a finite subcover, and therefore, E is not compact.
- (ii) If $y \in \overline{E} \setminus E$, then, similar to (i), we have that E contains points from infinitely many rows. Take one point from each of these rows and call the resulting set A . Then the open cover $\{T \setminus A\} \cup \{a\} : a \in A \cap E\}$ of E cannot be reduced to a finite subcover and hence E is not compact.

□

Proof of 1.1. Assume that there exists a surjective continuous map $f : C \rightarrow T$. We will show that this implies that (T, τ) is Hausdorff which contradicts Property (b) of Lemma 4.1.

Choose two distinct points $u, v \in T$. Since singletons are compact we have by Property (c) of Lemma 4.1 that the sets $\{u\}$ and $\{v\}$ are closed in T with respect to τ . Therefore, their pre-images under f are non-empty closed subsets of C and have disjoint open neighbourhoods $U(u)$, $U(v)$, as C is normal. The complements of these open neighbourhoods are compact subsets of C . Thus, their images under f are compact in T because f is continuous and they are closed because of property (3) of Lemma 4.1. Therefore,

$$V(u) := f(U(u)^c)^c \quad \text{and} \quad V(v) := f(U(v)^c)^c$$

are open neighbourhoods of u and v respectively. We claim that these sets are disjoint:

$$\begin{aligned} V(u) \cap V(v) &= f(U(u)^c)^c \cap f(U(v)^c)^c = (f(U(u)^c) \cup f(U(v)^c))^c \\ &= (f(U(u)^c \cup U(v)^c))^c \\ &= (f((U(u) \cap U(v))^c))^c = (f(\emptyset^c))^c = \emptyset. \end{aligned}$$

Hence we have separated the points u and v by open neighbourhoods. Since $u, v \in T$ were chosen arbitrarily, we conclude that (T, τ) is Hausdorff. □

Remark. In the above proof we demonstrated that there exists a countably infinite topological space which is not the continuous image of C . By adjoining a compact Hausdorff space, such as the unit interval, to the space (T, τ) , using the disjoint union topology, one obtains a compact non-Hausdorff space with cardinality equal to that of C . Arguments similar to those given above will then show that this space is not the continuous image of C . (See [11, 12] for more on disjoint union topologies.)

REFERENCES

- [1] P. S. Alexandroff, Über stetige Abbildungen kompakter Räume, *Math. Annalen* **96** (1927), 55–571.
- [2] G. Cantor, Grundlagen einer allgemeinen Mannigfaltigkeitslehre, *Math. Annalen* **21** (1883), 545–591.

- [3] K. J. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, second edition, John Wiley and Sons, 2003.
- [4] J. L. Fleron, A note on the history of the Cantor set and the Cantor function, *Mathematics Magazine*, **67** (1994), 136-140.
- [5] T. Hawkins, *Lebesgue's theory of integration: Its origins and development*, Chelsea Publishing Co., Madison, WI, 1975.
- [6] H. Hahn, Mengentheoretische Charakterisierung der stetigen Kurven, *S. B. Kaiserl. Akad. Wiss. Wien* **123** (1914), 2433-2487.
- [7] J. H. Manheim, *The genesis of point set topology*, Pergamon Press, Oxford, 1964.
- [8] L. R. Steen and J. A. Seebach (Jr.), *Counterexamples in Topology*, Dover Publications, Inc, New York, 1978.
- [9] H. J. S. Smith, On the integration of discontinuous functions, *Proc. London Math. Soc.* **6** (1875), 140-153; *Collected Mathematical Papers*, 25.
- [10] H. Sagan, *Space-Filling Curves*, Springer, New York, 1994.
- [11] W. A. Sutherland, *Introduction to Metric and Topological Spaces*, Oxford University Press, 1975.
- [12] S. Willard, *General Topology*, Addison-Wesley Publishing Company, 1970.